




Corporate Presentation

NASDAQ: CRIS



Cautionary Note Regarding Forward Looking Statements

This presentation contains certain forward-looking statements about Curis, Inc. (“we,” “us,” or the “Company”) within the meaning of the Private Securities Litigation Reform Act of 1995, as amended. Words such as “expect(s),” “believe(s),” “will,” “may,” “anticipate(s),” “focus(es),” “plans,” “mission,” “strategy,” “potential,” “estimate(s),” “intend,” “project,” “seek,” “should,” “would” and similar expressions are intended to identify forward-looking statements. Forward-looking statements are statements that are not historical facts, reflect management’s expectations as of the date of this presentation, and involve important risks and uncertainties. Forward-looking statements herein include, but are not limited to, statements with respect to the timing and results of future clinical and pre-clinical milestones; the timing of future preclinical studies and clinical trials and results of these studies and trials; the clinical and therapeutic potential of our drug candidates; the proposed focus on emavusertib and management’s ability to successfully achieve its goals. These forward-looking statements are based on our current expectations and may differ materially from actual results due to a variety of important factors including, without limitation, risks relating to: whether and when the U.S. Food and Drug Administration may remove the partial clinical hold on the combination therapy phase (Phase 1b) and the expansion phase (Phase 2a) of the Phase 1/2 TakeAim Leukemia trial, or may take further regulatory action with regard to this trial, whether any of our drug candidates will advance further in the clinical development process and whether and when, if at all, they will receive approval from the FDA or equivalent foreign regulatory agencies; whether historical preclinical results will be predictive of future clinical trial results; whether historical clinical trial results will be predictive of future trial results; whether any of our drug candidate discovery and development efforts will be successful; whether any of our drug candidates will be successfully marketed if approved; our ability to achieve the benefits contemplated by our collaboration agreements; management’s ability to successfully achieve its goals; the sufficiency of our cash resources; our ability to raise additional capital to fund our operations on terms acceptable to us or the use of proceeds of any offering of securities or other financing; general economic conditions; competition; and the other risk factors contained in our periodic and interim reports filed with the Securities and Exchange Commission which are available on the SEC website at www.sec.gov. You are cautioned not to place undue reliance on these forward-looking statements that speak only as of the date hereof, and we do not undertake any obligation to revise and disseminate forward-looking statements to reflect events or circumstances after the date hereof, or to reflect the occurrence of or non-occurrence of any events, except as required by law.

Corporate Overview


Summary

Investment Thesis	Curis develops novel cancer therapeutics in areas of significant unmet patient need <i>Cash runway into 2025 – \$98.7M as of Sept 30, 2022</i>
Lead Program	Emavusertib is positioned to become the cornerstone agent in heme malignancies • IRAK4-L is the most prevalent driver of disease in AML/MDS^{1,2} • IRAK4i has a synergistic effect when combined with ibrutinib in NHL
Market Opportunity	AML/MDS: 317K patients ³ <i>(current standard of care is HMA)</i> NHL/CLL: 1.8M patients ³ <i>(current standard of care is BTKi)</i>
2023 Milestones	Mid-Year: Full release of clinical hold on TakeAim Leukemia Mid-Year: Agreement with FDA on Recommended Phase 2 Dose 2 nd Half: Updated data from both the R/R Leukemia and R/R Lymphoma studies


1) Smith et al. Nat Cell Biol 2019; 2) Saygin, et al. J Hematol Oncol. 2017 Apr 18; 3) 2022 Prevalence Data DRG Clarivate

Pipeline

Curis develops first-in-class cancer therapeutics



¹ In April 2022, the U.S. Food and Drug Administration (“FDA”) placed the TakeAim Leukemia study on partial clinical hold. In August 2022, the FDA notified Curis that it may resume enrollment in the monotherapy dose finding phase of the study. The partial hold remains in place for the combination therapy and expansion phases of the study.

A large, circular, light blue microscopic image of a cell is centered behind the title. The cell has a textured, wavy surface and a darker blue, granular interior. The background of the slide is white with several out-of-focus, light blue, irregular shapes scattered around.


Emavusertib: Overview of Initial Clinical Data

Emavusertib Overview


Initial clinical data support development strategy for novel IRAK4 inhibitor

Initial Clinical Data in AML/MDS


(patients treated with monotherapy, grouped by targeted mutation status)




With Targeted Mutations Monotherapy



All Comers Combination



1 – in combination with ibrutinib
2 – in combination with azacitidine and/or venetoclax

A circular inset containing a blue-toned microscopic image of a cell, showing a textured, granular surface.


IRAK4 Biology and Mechanism of Action

Emavusertib Mechanism of Action

IRAK4 is a novel and important target across multiple heme malignancies


IRAK4 in AML/MDS

Spliceosome mutation drives overexpression of IRAK4-L (which causes constitutive activation of the myddosome)



IRAK4 in NHL/CLL

TLR Pathway is dependent upon IRAK4 for function (2nd pathway driving NF-κB overactivity)



1) Guillaumot et al. Nat Cell Biol 2019; 2) Smith et al. Nat Cell Biol 2019; 3) American Cancer Society, Cancer Facts & Figures 2020; 4) Leukemia & Lymphoma Society, Facts and Statistics Overview; 5) IMBRUVICA Package Insert. Rev 08/2018


Emavusertib Unique Molecular Fingerprint

Targeted design specifically engineered to hit key oncogenic targets

The NCI has selected emavusertib for both clinical and non-clinical studies of IRAK4 inhibition in oncology

NCI Selection for IRAK4

Emavusertib
Kinase Interaction Map



% Inhibition at 0.1 nM
Illustration reproduced courtesy of Cell Signaling Technology

Emavusertib
Binding Affinity

Target	K _d nM
IRAK1	12,000
IRAK2	>20,000
IRAK3	8,500
IRAK4	23
DYRK1A	25
FLT3 wt	31
FLT3 (D835H)	5
FLT3 (D835V)	44
FLT3 (D835Y)	3
FLT3 (ITD)	8
FLT3 (K663Q)	47
FLT3 (N841I)	16
Haspin (GSG2)	32
CLK1	10
CLK2	20
CLK3	>20,000
CLK4	14
TrkA	130

DiscoverX Kinase Panel
(378 kinases screened)


Binds with high affinity to IRAK4

Dual targeting of IRAK4 and FLT3 confers potential efficacy advantage vs. other IRAK4 inhibitors

Emavusertib Preclinical Data


Clear anti-cancer activity suggests broad potential across heme malignancies

AML/MDS Monotherapy




emavusertib demonstrates monotherapy activity in patient-derived xenografts¹

AML/MDS Combination Therapy




emavusertib demonstrates synergy with both azacitidine and venetoclax in THP-1 model²

NHL/CLL Combination Therapy



emavusertib demonstrates synergy with ib Brutinib in OCI-Ly10 model³

A large, circular, light blue microscopic image of a cell is centered in the background. The cell has a textured, wavy surface and a darker blue, granular interior. A white horizontal band is superimposed over the center of the cell, containing the text.


Emavusertib in Leukemia (AML/MDS)

Emavusertib in AML/MDS with FLT3 Mutation


Preclinical data suggest potential to address FLT3 population, as well as spliceosome population

“Concomitant targeting of IRAK1 or IRAK4, alongside FLT3, is the most effective means to overcome the adaptive resistance incurred when targeting FLT3”


IRAK activity increases after treatment with FLT3i



IRAK/FLT3 combination is synergistically cytotoxic




Mice die if treated FLT3i alone, but survive if treated with IRAK/FLT3 combination



Emavusertib Initial Clinical Data

Potential for clear regulatory path in two target populations: AML FLT3 and AML Spliceosome




	CR/CRh Rate	
FLT3 AML	1/3	(33%)
Spliceosome AML	2/5	(40%)

- All patients were R/R, previously treated with HMA
 - Prognosis for this population is very poor; mOS is 2-4 months¹
- Full approval has been granted for several AML drugs with a single-arm study using CR/CRh rate as primary endpoint
 - 21% CR/CRh rate for gilteritinib
 - 23% CR/CRh rate for enasidenib

1) Maiti et al. Haematologica 2021

Emavusertib Initial Clinical Data

Clear unmet need in R/R hrMDS – there are no approved therapies



Spliceosome MDS

ORR

4/7 (57%)


- All patients on study were R/R, previously treated with HMA
 - Prognosis for this population is very poor; mOS is 4-6 months¹
 - There are no approved therapies for patients R/R to HMA
- U2AF1/SF3B1 are the most prominent mutations in MDS
 - 30% of MDS patients have one of these two spliceosome mutations²

Emavusertib in AML/MDS

Genomic data provide clear rationale for monotherapy & combination

IRAK4-L is the most prevalent disease driver in AML/MDS^{1,2}

<u>Disease Driver</u>	<u>% of Patient Population</u>
IRAK4-L	> 50% ¹
FLT3	> 25% ²
TET2	10-20% ³
IDH2	9-13% ⁴
IDH1	6-10% ⁴
CEBPA	~10% ³



- IRAK4 activation stimulates NF- κ B and an array of anti-apoptotic factors
- emavusertib enhances the anti-cancer efficacy of azacitidine and venetoclax in preclinical models

Emavusertib in AML/MDS

Clinical data support strategy for monotherapy & combination

In patients with a targeted mutation:

- Enhanced monotherapy efficacy in a genetically-defined population suggests potential for clear path to first NDA submission


In patients without a targeted mutation:

- Initial data show broad anti-cancer activity in non-targeted patients, suggesting emavusertib could enhance the efficacy of other therapies when used in combination

***Spliceosome mutations can be used
to identify patients with higher levels of IRAK4-L expression***

- *U2AF1 and SF3B1 mutations cause overexpression of IRAK4-L¹*
- *Genetic screening enables the utilization of existing gene panels to identify candidates for monotherapy*

Initial Clinical Data in AML/MDS
(patients treated with monotherapy, grouped by targeted mutation status)



1) Guillaumot et al. Nat Cell Biol 2019

Emavusertib in Leukemia (AML/MDS)

The ideal candidate to address the two largest genetically-defined populations in AML/MDS

Emavusertib addresses the two largest genetically-defined populations in AML/MDS:

- 1) IRAK4-L (>50% of population)¹
- 2) FLT3 (>25% of population)²


- Clear single-agent activity
- Genetically-defined patient population
- Dual targeting of IRAK4 and FLT3 may prevent adaptive resistance³ observed with current FLT3 inhibitors



Next Steps

TakeAim Leukemia Study

- *Monotherapy: targeted patients (spliceosome, FLT3)*
- *Combination: all other patients*

A large, circular, light blue microscopic image of a cell is centered in the background. The cell has a textured, wavy surface and a darker blue nucleus. A white horizontal bar is superimposed over the center of the cell, containing the text.


Emavusertib in Lymphoma (NHL/CLL)

Emavusertib in B Cell Cancers

Combination therapy provides complimentary inhibition of two pathways that drive NF- κ B

NF κ B Biology: Two Pathways Drive B Cell Cancers


*BCR and TLR Pathways independently drive NF- κ B overactivity
(and NF- κ B drives B Cell Cancers)*



1) IMBRUVICA Package Insert. Rev 08/2018

Clinical Strategy: Use Combination Therapy


*In preclinical testing, blocking both IRAK4 and BTK
drove tumor reduction better than blocking either one alone*




2) Booher et al. Waldenström Roadmap Symposium 2019

Emavusertib Initial Clinical Data

Majority of patients achieved decreases in tumor burden, including complete responses



13 total patients enrolled
9 patients were evaluable for tumor burden
4 patients did not receive pre/post tumor assessments



Emavusertib in Lymphoma (NHL/CLL)

The ideal candidate to combine with BTKi to maximize downregulation of NF- κ B

- Patients are currently treated with BTKi because it downregulates NF- κ B
- Two pathways drive NF- κ B:
 - 1) BCR Pathway: addressed by blocking BTK
 - 2) TLR Pathway: addressed by blocking IRAK4
- Initial clinical data suggest blocking both pathways may overcome resistance to ibrutinib



Next Steps

TakeAim Lymphoma Study

- Targeted Patients: MZL, pCNSL
- All Comers: patients resistant to BTKi

Corporate Overview

Summary

Investment Thesis	Curis develops novel cancer therapeutics in areas of significant unmet patient need <i>Cash runway into 2025 – \$98.7M as of Sept 30, 2022</i>
Lead Program	Emavusertib is positioned to become the cornerstone agent in heme malignancies • IRAK4-L is the most prevalent driver of disease in AML/MDS^{1,2} • IRAK4i has a synergistic effect when combined with ibrutinib in NHL
Market Opportunity	AML/MDS: 317K patients ³ <i>(current standard of care is HMA)</i> NHL/CLL: 1.8M patients ³ <i>(current standard of care is BTKi)</i>
2023 Milestones	Mid-Year: Full release of clinical hold on TakeAim Leukemia Mid-Year: Agreement with FDA on Recommended Phase 2 Dose 2 nd Half: Updated data from both the R/R Leukemia and R/R Lymphoma studies

1) Smith et al. Nat Cell Biol 2019; 2) Saygin, et al. J Hematol Oncol. 2017 Apr 18; 3) 2022 Prevalence Data DRG Clarivate



End of Corporate Presentation

NASDAQ: CRIS

